

Distribution of extreme waves by simulations.

Hervé Socquet-Juglard and Kristian Dysthe, University of Bergen

Karsten Trulsen, University of Oslo

Harald E.Krogstad and Jingdong Liu, NTNU, Trondheim

Abstract

Several ideas for physical mechanisms that can produce large waves have been suggested as reasons for rogue- or freak waves (see e.g. Dysthe 2000, Kharif & Pelinovsky 2004). It seems, however, that these mechanisms need special preparations or coherence to work. It is therefore difficult to see how they can occur spontaneously during a storm on the open ocean.

To investigate the possibility that nonlinear wave interactions may influence the spontaneous occurrence of extreme waves, we have done fairly large scale 3D simulations using higher order NLS type equations (Dysthe 1979). Our "numerical" ocean contains approximately 10^4 waves at any time, and has a time horizon of 150 typical wave periods $2\pi/\omega_p$. As has been reported elsewhere (Dysthe et.al. 2003) a characteristic evolution is seen where the initial wave spectrum "relaxes" towards a more steady form. This process is most prominent for very narrow spectra, and takes place on the Benjamin-Feir timescale $(\omega_p s^2)^{-1}$ where s is a typical steepness. Such spectral change due to the Benjamin-Feir type instability has been linked to increased occurrence of extreme waves (Onorato et.al. 2000, Mori & Yasuda 2000, Janssen 2003). This connection is investigated using our simulations.

We are concerned with the distributions of surface elevation and crest height. For relatively short crested waves the mentioned spectral evolution does not seem to influence these distributions significantly. We find that up to 5 standard deviations the theoretical distributions found by Tayfun (1980) seems to fit the simulated data very well. For a very narrow initial angular distribution, however, we see a significant increase in the occurrence of extreme waves during the spectral relaxation, in qualitative agreement with laboratory experiments done by Onorato et.al. (2004).

Acknowledgements

This research has been supported by the BeMatA programme of the Research Council of Norway, and also by Statoil and Hydro.

References

DYSTHE, K.B. 2000 Modelling a "rogue wave"-speculations or a realistic possibility? *Proceedings of the workshop Rogue Waves 2000 IFREMER*, pp. 255-264 .

KHARIF, C. & PELINOVSKY, E. 2004 Physical mechanisms of the rogue wave phenomenon. *European Journal of mechanics B/Fluids*. **22**, pp. 603-634.

DYSTHE, K.B. 1979 Note on a modification to the nonlinear Schrödinger equation for application to deep water waves *Proc. R. Soc. Lond. A* **369**, pp. 105–114.

DYSTHE K.B., TRULSEN, K., KROGSTAD, H.E. & SOCQUET-JUGLARD, H. 2003 Evolution of a narrow-band spectrum of random surface gravity waves. *J. Fluid Mech.* **478**, pp. 1-10.

MORI, N and YASUDA, T. 2000 Effects of high-order nonlinear wave-wave interactions on gravity waves. *Proceedings of the workshop Rogue Waves 2000 IFREMER* pp. 229-244.

ONORATO, M., OSBORNE, A.R., SERIO, M. and DAMIANI, T. 2000 Occurrence of freak waves from envelope equations in random ocean wave simulations. *Proceedings of the workshop Rogue Waves 2000 IFREMER*. pp. 181-191.

JANSSEN, P.A.E.M. 2003 Nonlinear four wave interaction and freak waves. *J.Phys.Oceanogr.* **33**, pp. 863-884.

TAYFUN, M.A. 1980 Narrow-band nonlinear sea waves. *J. Geophys. Res.* **85**, pp. 1548-1552.

ONORATO, M., OSBORNE, A.R., SERIO, M., CAVALERI, L., BRANDINI, C. and STANSBERG, C.T. 2004 Extreme waves and modulational instability: wave flume experiments on irregular waves. *Submitted to J. Fluid Mech.*